



# New Long-Term Care Residents Who Potentially Could Have Been Cared for at Home

## Appendices



Production of this document is made possible by financial contributions from Health Canada and provincial and territorial governments. The views expressed herein do

---

---

# Table of contents


Appendix A: Describing home care clients — numerator definition . . . . .	4
Appendix B: Outcome scales in RAI-MDS 2.0 and interRAI LTCF . . . . .	5
Appendix C: Text alternative for flowchart . . . . .	6

# Appendix A: Describing home care clients — numerator definition

Methodological approach: To identify the characteristics of home care clients living well at home through home care programs.

This indicator identifies newly admitted long-term care residents who have a clinical profile similar to that of clients cared for at home with formal supports in place. Examples of formal home care supports include help with daily tasks such as bathing, dressing, eating and/or toileting.

The clinical characteristics of long-term care residents in the flowchart below represent the **numerator** of this indicator.



## Appendix B: Outcome scales in RAI-MDS 2.0 and interRAI LTCF

Outcome scale	Description	RAI-MDS 2.0 assessment items	interRAI LTCF assessment items	Score range
Activities of Daily Living (ADL) Hierarchy Scale	This scale reflects the disablement process by grouping ADL performance levels into discrete stages of loss (early loss: personal hygiene; middle loss: toileting and locomotion; and late loss: eating).	4 ADL Self-Performance Hierarchy Scale items: <ul style="list-style-type: none"> <li>• Personal Hygiene (G1jA)</li> <li>• Toilet Use (G1iA)</li> <li>• Locomotion (G1eA)</li> <li>• Eating (G1hA)</li> </ul>	<ul style="list-style-type: none"> <li>• Personal hygiene (G1b)</li> <li>• Locomotion (G1f)</li> <li>• Toilet use (G1h)</li> <li>• Eating (G1j)</li> </ul>	0–6 Higher scores indicate greater decline (progressive)

## Appendix C: Text alternative for flowchart

To identify the characteristics of home care clients who are living well at home through home care programs, we first look at home care clients who were assessed in a private home, a condominium, an apartment or an assisted-living setting between January 1, 2014, and March 31, 2019. From that group, we then identify home care clients who were on service for at least 180 days at the time of the assessment and who remained on service for at least 180 days after the assessment. The client and caregiver must indicate that they do not believe the client will be better off elsewhere. From this group, we then identify clients who are below the 80th percentile for home care service hours (i.e., received less than 750 minutes per week).

We then look at the characteristics of these home care clients who, over a 5-year period, are living well in the community with formal supports: no to mild cognitive impairment (Cognitive Performance Scale = 0–2); at a maximum, requires limited assistance in activities of daily living (ADLs) (ADL Hierarchy Scale = 0–2); no falls in the past 90 days; not physically or verbally abusive; and no wandering.

To identify newly admitted long-term care residents who potentially could have been cared for at home, we

---