

How Canada Compares

Results From the Commonwealth Fund's 2019 International Health Policy Survey of Primary Care Physicians

Accessible Report

January 2020

n stitute
f forn a
Institut canadi- nté

Production of this document is made possible by financial contributions from Health Canada and provincial and territorial governments. The views expressed herein do not necessarily represent the views of Health Canada or any provincial or territorial government.

Unless otherwise indicated, this product uses data provided by Canada's provinces and territories.

All rights reserved.

The contents of this publication may be reproduced unaltered, in whole or in part and by any means, solely for non-commercial purposes, provided that the Canadian Institute for Health Information is properly and fully acknowledged as the copyright owner. Any reproduction or use of this publication or its contents for any commercial purpose requires the prior written authorization of the Canadian Institute for Health Information. Reproduction or use that suggests endorsement by, or affiliation with, the Canadian Institute for Health Information is prohibited.

For permission or information, please contact CIHI:

Canadian Institute for Health Information
495 Richmond Road, Suite 600
Ottawa, Ontario K2A 4H6
Phone: 613-241-7860
Fax: 613-241-8120
cihi.ca
copyright@cihi.ca

ISBN 978-1-77109-888-5

© 2020 Canadian Institute for Health Information

How to cite this document:
Canadian Institute for Health Information. *How Canada Compares: Results From the Commonwealth Fund's 2019 International Health Policy Survey of Primary Care Physicians — Accessible Report*. Ottawa, ON: CIHI; 2020.

Cette publication est aussi disponible en français sous le titre *Résultats du Canada : Enquête internationale de 2019 du Fonds du Commonwealth sur les politiques de santé auprès des médecins de soins primaires — rapport en format accessible*.
ISBN 978-1-77109-889-2

Table of contents

Executive summary 4

Acknowledgements 6

About this report 7

Reporting framework 8

3UR¿OH RI SULPDU\ FDUH SK\VLFLDQV DQG

Patient-centered care

- The majority of Canadian primary care physicians feel well prepared to care for patients with chronic conditions (82%). In contrast, fewer Canadian primary care physicians feel prepared to care for patients with specialized needs, particularly dementia (40%), palliative care (36%) and substance use (19%). In Canada, 13% of primary care physicians reported feeling well prepared to care for patients requesting medical assistance in dying.

Coordination within the health system with social services

- Fewer Canadian primary care physicians communicate with home care providers about their patients' needs (24%) compared with the CMWF average (31%). However, about the same proportion of Canadian physicians receives updates about their patients (36%) as the CMWF average (37%).
- Although many Canadian primary care physicians (60%) screen their patients for social needs, fewer frequently coordinate care with social services (43%). One of the biggest challenges is LQDGHTXDWH VWD QJ WR PDNH UHIHUUDOV DQG FRRUGLQDWH across jurisdictions.
- 65% of Canadian primary care physicians think that better integration of primary care with hospitals, mental health services and community-based social services is the top priority in improving quality of care and patient access.

Coordinating information technologies

- More Canadian primary care physicians were using electronic medical records (EMRs) in 2019 (86%) than in 2015 (73%), but this was still lower than the CMWF average (93%).
-)HZHU &DQDGLDQ SULPDU\ FDUH SUDFWLFHV R HU WKHLU SDW their patient visit summaries online (Canada: 5%; CMWF: 26%) and request prescription renewals online (Canada: 10%; CMWF: 52%). Similarly, compared with the CMWF average, fewer Canadian primary care practices can exchange information electronically with doctors outside their practice, including patient clinical summaries (Canada: 25%; CMWF: 63%), laboratory and diagnostic test results (Canada: 36%; CMWF: 65%) and lists of medications taken by their patients (Canada: 33%; CMWF: 62%).
- Compared with the CMWF average, fewer Canadian primary care physicians review their performance on clinical outcomes (Canada: 34%; CMWF: 60%), patients' hospital admissions (Canada: 25%; CMWF: 32%), prescribing practices (Canada: 26%; CMWF: 58%), surveys of patient satisfaction and experiences with care (Canada: 17%; CMWF: 38%) and surveys of patient-reported outcome measures (Canada: 8%; CMWF: 22%).

Acknowledgements

The Commonwealth Fund provided core support, with cofunding from the German Federal Ministry of Health and the German Institute for Quality Assurance and Transparency in Healthcare (Germany); the Haute autorité de santé, Caisse nationale de l'assurance maladie and the Directorate for Research, Evaluation, Studies, and Statistics of the French Ministry of Health (France); the Dutch Ministry of Health, Welfare, and Sport, and Radboud University Medical Center (the Netherlands); the Norwegian Institute of Public Health (Norway); the Swedish Agency for Health and Care Services Analysis (Sweden); and the Swiss Federal 2 F H R I 3 X E O L F + H D O W K 6 Z L W] H U O D Q G $ G G L W L R Q D O V X S S R U provided by the New South Wales Agency for Clinical Innovation and the Department of Health and Human Services (Australia); the Canadian Institute for Health Information, Canada Health Infoway, the ministère de la Santé et des Services sociaux du Québec and Health Quality Ontario (Canada); and the Health Foundation of the United Kingdom (U.K.).

Within Canada, funding for an expanded Canadian sample was provided by CIHI, Canada Health Infoway, the ministère de la Santé et des Services sociaux du Québec and Health Quality Ontario.

CIHI would like to acknowledge and thank the many individuals who assisted with the development of this report, including our Expert Advisory Group:

- Dr. Mike Benigeri, Consultant
- Caroline Boucher, Strategic Planning Advisor, ministère de la Santé et des Services sociaux du Québec
- Dr. Bidénam Kambia-Chopin, 3 H U I R U P D Q F H $ S S U D L V DCommissaire à la F H 2 F H U santé et au bien-être du Québec
- Dr. Susan Chatwood, 6 F L H Q W L ¿ F ' L U H F W R U , Q V W L W X W H I R U & L U F X P S
- Tara S. Chauhan, Senior Advisor, Canadian Medical Association
- Dr. Gail Dobell, Interim Vice President, Health System Performance, Health Quality Ontario
- Annette McKinnon, patient representative
- Michelina Mancuso, Executive Director, Performance Measurement, New Brunswick Health Council
- Dr. Alex Singer, Director, Manitoba Primary Care Research Network
- Sukirtha Tharmalingam, Manager, Evaluation Methods, Canada Health Infoway

Please note that the analyses and conclusions in the present document do not necessarily U H À H F W W K R V H R I W K H L Q G L Y L G X D O V R U R U J D Q L] D W L R Q V P H Q W

H

Sample sizes in some provinces are much smaller than in others and have wider margins of error. For this reason, provinces may have the same or different results, e.g., one result might be the same as or different from the average. The most robust samples are in Quebec and Ontario because of the additional funding from those provinces.

To assess the reliability of the results, the coefficient of variation (CV) is the standard error of an estimate expressed as a percentage of the estimate and is a measure of sampling error. Estimates with a CV less than 16.6% are considered reliable for general use. Estimates with a CV between 16.6% and 33.3% are considered to have high levels of error, and caution should be exercised when using results with such CVs. Estimates with a CV higher than 33.3% are considered unreliable and are suppressed, as indicated with an em dash (—). These quality level guidelines are consistent with those used at Statistics Canada.[1]

Some of the questions were part of the Canadian survey only and were not asked in other countries. International comparison is therefore not possible. Findings for these questions are

Profile of primary care physicians and their practices

Key findings

- Canadian physicians see a median of 100 patients a week, similar to the 11-country average of medians (99 patients a week). 55% of Canadian primary care physicians spend 15 to less than 25 minutes with each patient, similar to the CMWF average (54%). Most Canadian primary care practices (61%) are not accepting new patients.
-

Differences between solo and group physician practices

Table 3

Table 4 Median number of patients seen during a typical workweek, by jurisdiction

Jurisdiction	Median patients seen
N.L.	125
P.E.I.	100
N.S.	112
N.B.	110
Que.	70
Ont.	100
Man.	100

Canadian physicians work similar hours as CMWF average

Table 5 Proportion of primary care physicians by the number of hours they typically work each week in their medical practice, by country

Canadian primary care physicians spend similar amount of time with patients as CMWF average

Table 6 Proportion of primary care physicians by the average amount of time they are able to spend with a patient during a routine office visit, by country

Country	Less than 15 minutes	15 to less than 20 minutes	20 minutes or more
Sweden	2	44	54
Norway	3	86	12
Switzerland	6	66	27
France	8	82	10
New Zealand	10	87	3
United States	16	63	21
Australia	25	70	5
Canada	28	55	18
CMWF Average	32	54	14
Netherlands	85	15	0
Germany	85	12	2
United Kingdom	86	14	0

&RPSDUHG ZLWK WKHUH ZHUH QR VLJQL¿FDQW FKDQJHV LQ SULPDU\ FDUH SK\VLFLDQV ZHUH DEOH WR VSHQG ZLWK SDWLHQ

Physician satisfaction by time spent with patients

Table 7 Proportion of Canadian primary care physicians who are extremely, very or moderately satisfied with the time they can spend with patients, by the amount of time they are able to spend with patients during routine office visits

Amount of time	Extremely satisfied
Less than 15 minutes	53
15 to less than 20 minutes	72
20 minutes or more	82

57% of Canadians feel that their regular doctor always spends enough time with them when they need care or treatment.[2]

Provincial and territorial snapshot: Time

Table 8 Proportion of primary care physicians by the number of hours they typically work each week in their medical practice, by jurisdiction [d]

Table 8b Proportion of primary care physicians by the average amount of time they are able to spend with a patient during a routine office visit, by jurisdiction [d]

Table 9 Of those who have the capacity and are accepting new patients, proportion of primary care physicians who use the following strategies,† Canada

Strategy	Percentage
Use a waiting list maintained by the clinic	46
Use the public waiting list for patients maintained by the government	36
Put them in the government's system to be seen	33
Use both strategies if I can	59

Tbe 10b

Table 11b Trend over time, proportion of primary care physicians who feel extremely or very satisfied with their job as a primary care physician

Country	2015	2019
Canada	27	46[b]
Commonwealth Average	35	45

Notes
b: Same as average.
Lower numbers are desirable.

Provincial and territorial snapshot: Satisfaction and stress

Table 12 Proportion of primary care physicians who are extremely, very or moderately satisfied with aspects of medical practice, by jurisdiction

Jurisdiction													

Table 12 Proportion of primary care physicians who feel extremely or very stressed with their job as a primary care physician,† by jurisdiction

Jurisdiction	N.L.	P.E.I.	N.S.	N.B.	Que.	Ont.	Man.	Sask.	Alt.	B.C.	Terr.	Can.	CMWF Avg.
Percentage	46[b]	[b]	61[c]	42[b]	34[a]	55[c]	43[b]	45[b]	40[b]	48[b]	49[d]	46[b]	45

Notes

7KH FRH FLHQW RI YDULDWLRQ &9 LV EHWZHHQ DQG XVH ZLWK FDXWLRQ

† Lower results are more desirable.

a: Above average.

b: Same as average.

c: Below average.

d: Non-directional or no statistical tests were performed to compare with the CMWF average (e.g., territories).

More Canadian primary care physicians offer weeknight and weekend appointments than CMWF average

Table 13

Proportion of primary care physicians whose practice offers appointments on weekends (i.e., Saturday or Sunday) to see patients, by country

Fewer Canadian primary care physicians have after-hours arrangements

Table 15: Proportion of primary care physicians whose practice has an arrangement, either internally or with another practice, where patients can be seen by a doctor or nurse when the practice is closed (e.g., after hours), by country

Country	Percentage
Germany	96
Switzerland	92
Netherlands	91
Australia	90
United Kingdom	84
Sweden	77
France	75
New Zealand	75
Italy	69
Norway	56
Canada	49[c]
United States	45

Table 15b Of those who have arrangements, proportion of primary care physicians who by receive notifications that their patients have been seen for after-hours care, by country

Country	Percentage
Australia	98
New Zealand	79
United Kingdom	77
Canada[c]	45
Austria	43
Italy	43
United States	37
Germany	37
Switzerland	29
France	28[c]
Sweden	11
Norway	8

Note
c: Below average.

In 2015, Canadian primary care physicians reported similar results for having arrangements when the practice is closed. Canada's place among other countries was similar in both 2015 and 2019.[4]

Fewer Canadian primary care physicians offer access to patients electronically compared with CMWF average

Table 17b: Proportion of primary care physicians whose practice offers patients the option to request appointments online (not including email), by country

Country	Percentage
United Kingdom	91
Sweden	87
Norway	83

Provincial and territorial snapshot: Electronic access

Table 18 Proportion of primary care physicians whose practice offers patients electronic access options, by jurisdiction

Jurisdiction	N.L.	P.E.I.	N.S.	N.B.	Que.	Ont.	Man.	Sask.	Alt.	B.C.	Terr.	Can.	CMWF Avg.
Connected with their practice via secure website to email questions or concerns	[c]	—	25[c]	[c]	17[c]	30[c]	[c]	[c]	[c]	24[c]	—	23[c]	65
Requests appointments online (including email)	—	—	[c]	[c]	32[c]	19[c]	27[c]	[c]	[c]	27[c]	0[d]	22[c]	56

Notes

7KH FRH FLHQW RI YDULDWLRQ &9 LV EHWZHHQ DQG XVH ZLWK FDXWLRQ

— Data is suppressed due to extreme sampling variability (CV>33.3%).

Table 19 Proportion of primary care physicians who responded that they or other health care professionals in their practice frequently make home visits, by jurisdiction

Jurisdiction	Percentage
N.L.	33[c]
P.E.I.	—
N.S.	[c]
N.B.	[c]
Que.	25[c]
Ont.	18[c]
Man.	[c]
Sask.	[c]
Alt.	[c]
B.C.	[c]
Ter.	[d]

Notes

Patient-centred care

Key findings

- The majority of Canadian primary care physicians feel well prepared to care for patients

Provincial and territorial snapshot: Specialized needs

Table 2 Proportion of primary care physicians whose practice is all or most of the time, with respect to having sufficient skills and experience, to manage care for patients with specialized needs, by type of need and jurisdiction

Fewer Canadian primary care physicians use personnel to care for patients with chronic conditions compared with CMWF average

Table 2 Proportion of primary care physicians who use personnel, such as nurses or case managers, to monitor and manage care for patients with

Table 2 Proportion of primary care physicians who use personnel, such as nurses or case managers,

Management of chronic conditions

Table 2 Proportion of primary care physicians who responded that they or other health care professionals in their practice usually or often provide care for patients with chronic conditions in the following ways

Chronic condition management activities	Canada	CMWF Average
Develop plans with patients they can rely on in the event of a crisis	74[a]	69
Provide patients with written instructions		4

Most Canadian primary care physicians have

Provincial and territorial snapshot: Palliative and end-of-life care

Table 2 End-of-life care, by proportion of primary care physicians and jurisdiction

Few Canadian primary care physicians well prepared to manage patients requesting medical assistance in dying (Canada only)

Table 2 Proportion of primary care physicians whose practice is [illegible], with respect to having sufficient skills and experience, to manage care for patients requesting medical assistance in dying, by jurisdiction [d]

Jurisdiction	Percentage
N.	
P.E.I.	—
N	
N	24
Que.	12
Ont	12
Man	
Sask	
Alt	
B.	
Ter	
Can	13

Notes

[illegible]

— Data is suppressed due to extreme sampling variability (CV>33.3%).

d: Non-directional or no statistical tests were performed to compare with the CMWF average (e.g., territories).

The Canadian average represents the average experience of Canadians (as opposed to the mean of provincial and territorial results).

Dying today

Medical assistance in dying is not considered part of palliative care, but it is an end-of-life option for Canadians who meet the legal criteria.

Between December 10, 2015 (when the law was enacted) and October 31, 2018, there were 6,749 medically assisted deaths in Canada.[9]

Coordination within the health system

Key findings

- While more Canadian primary care physicians send information to specialists (90%) compared with the CMWF average (85%), timely transfer of information back from specialists is lower in Canada (17% versus 21%).
- 7KH SURSRUWLRQV RI &DQDGLDQ SULPDU\ FDUH SK\VLFLDQV Z patients had visited the emergency department (49%) and had been hospitalized (54%) are similar to the CMWF averages (51% and 55%, respectively).
- Fewer Canadian primary care physicians communicate with home care providers about their patients' needs (24%) compared with the CMWF average (31%). However, about the same proportion of Canadian physicians receives updates about their patients (36% versus 37%).

Two-way communication between primary care physicians and specialists

Table 2 Proportion of primary care physicians who responded that they usually do the following when their patients have been referred to a specialist

Activities	Canada	CMWF average
Send the specialist the reason for the referral	90[a]	85
Receive from the specialist findings, changes to the patient's medication or treatment plan	59[b]	58
Receive report from the specialist within 1 week of service	17[c]	21

Notes
a: Above average.
b: Same as average.
c: Below average.

In 2016, 13% of Canadians reported that their specialist did not have basic information or test results from their regular doctor about the reason for their visit,[2] and 21% reported that their regular doctor did not seem informed and up to date about the care they received from the specialist.[2]

Provincial and territorial snapshot: Communication with specialists

Table 9 Proportion of primary care physicians who responded that they usually do the following when their patients have been referred to a specialist, by jurisdiction

A quarter of Canadian primary care physicians communicate with home-based nursing care about their patients' needs

Table 3 For patients who receive home-based nursing care, proportion of primary care physicians who say they communicate with home-based nursing care providers about their patients' needs and the services to be provided,† by country

Table 3.4

For patients who receive home-based nursing care, proportion of primary care physicians who say they are advised by the home-based nursing care providers of a relevant change in their patients' condition or health status,† by country

Country	Percentage
Switzerland	47
Sweden	46
France	45
Norway	43
United Kingdom	43
United States	42
Germany	38
Commonwealth Fund Average	37
Canada	36[b]
Australia	28
New Zealand	23
Netherlands	21

Notes

† Excludes respondents who answered "does not apply."

b: Same as average.

c: Below average.

Provincial and territorial snapshot: Communication with home-based nursing care

Table 5 For patients who receive home-based nursing care, proportion of primary care physicians who say they have the following communications with home-based nursing care providers, by jurisdiction

Table 8

Proportion of primary care physicians who responded that they or other health care professionals in their practice rarely or never coordinate care with social services or other community providers, by country

Country	

Provincial and territorial snapshot: Assessing social needs and coordinating social services

Table 3.1 Proportion of primary care physicians who usually or often screen or assess patients for at least one type of social need, by jurisdiction

Jurisdiction	N.L.	P.E.I.	N.S.	N.B.	Que.	Ont.	Man.	Sask.	Alt.	B.C.	Terr.	Can.	CMWG
Percentage													

Coordination using information technologies

Key findings

- More Canadian primary care physicians were using EMRs in 2019 (86%) than in 2015

Canadian primary care physicians catching up in use of EMRs

Table 4 Proportion of primary care physicians who use EMRs in their practice (not including billing systems), by country

Country	Percentage
New Zealand	100
Norway	100
United Kingdom	100
Netherlands	99
Sweden	99
Australia	97
CMWF Average	93

Provincial and territorial snapshot: EMR use

Table 4 Proportion of primary care physicians who use EMRs in their practice (not including billing systems), by jurisdiction

Jurisdiction	Percentage
N.L.	61[c]
P.E.I.	[c]
N.S.	86[c]
N.B.	61[c]
Que.	84[c]
Ont.	89[c]
Man.	88[b]
Sask.	91[b]
Alt.	92[b]
B.C.	90[b]
Terr.	96[d]
Can.	86[c]
CMWF Avg.	93

Notes

7KH FRH FLHQW RI YDULDWLRQ &9 LV EHWZHHQ DQG XVH ZLWK FDXWLRQ

b: Same as average.

c: Below average.

d: Non-directional or no statistical tests were performed to compare with the CMWF average (e.g., territories).

The CMWF average was calculated by adding the results from the 11 countries and dividing by

Electronic access to regional, provincial or territorial information systems (Canada only)

Table 4

Proportion of primary care physicians who have electronic access to any regional (e.g., hospital/ hospital network), provincial or territorial information systems where they can see patient information that is from outside their practice, by jurisdiction [d]

Jurisdiction	Percentage
N.L.	93
P.E.I.	91
N.S.	57
N.B.	94
Que.	76
Ont.	64
Man.	80
Sask.	92
Alta.	95
B.C.	57
Terr.	80
Can.	73

Notes
d: Non-directional or no statistical tests were performed to compare with the CMWF average (e.g., territories).
The Canadian average represents the average experience of Canadians (as opposed to the mean of provincial and territorial results).

Less than half of Canadian primary care physicians routinely use EMRs to support quality-of-care decisions

Table 4 Trend over time, proportion of primary care practices that do the following tasks routinely using a computerized system (e.g., EMR)

Fewer Canadian primary care physicians

Provincial and territorial snapshot: Performance review

Table 50 Proportion of primary care physicians whose practice receives and reviews data on the following aspects of their patients' care, [illegible] or [illegible], by jurisdiction

Type of data	N.L.	P.E.I.	N.S.	N.B.	Que.	Ont.	Man.	Sask.	Alta.	B.C.	Terr.	Can.	CMWF avg.
Clinical tests	23[c]	—	29[c]	50[c]	14[c]	40[c]	37[c]	31[c]	51[b]	37[c]	[d]	34[c]	60
Patients' hospital admissions or emergency department visits	[c]	[b]	[c]	40[b]	19[c]	30[b]	[c]	26[b]	29[b]	[c]	—	25[c]	32
Prescribing practices	24[c]	—	[c]	22[c]	10[c]	29[c]	30[c]	[c]	59[b]				

Perception of health system

Table 51b

Table 52 Proportion of primary care physicians who rate the overall performance of the health care system as Poor or Very Poor, by jurisdiction

/Type/Page/MediaBox[0 0 792 612]/Parent 2 0 R/Resources<</ProcSet[/PDF/Text]/ExtGState 3 0 R/Font<</F1 4 0 R/F2 5 0 R>>>>/Contents 6 0 R/Tabs/S/StructParents 0>>endobj 7 0 obj<</Type/Page/MediaBox[0 0 792 612]/Parent 2 0 R/Resources<</ProcSet[/PDF/Text]/ExtGState 3 0 R/Font<</F1 4 0 R/F2 5 0 R>>>>/Contents 8 0 R/Tabs/S/StructParents 1>>endobj 9 0 obj<</Type/Page/MediaBox[0 0 792 612]/Parent 2 0 R/Resources<</ProcSet[/PDF/Text]/ExtGState 3 0 R/Font<</F1 4 0 R/F2 5 0 R>>>>/Contents 10 0 R/Tabs/S/StructParents 2>>endobj 11 0 obj<</Type/Page/MediaBox[0 0 792 612]/Parent 2 0 R/Resources<</ProcSet[/PDF/Text]/ExtGState 3 0 R/Font<</F1 4 0 R/F2 5 0 R>>>>/Contents

Provincial and territorial snapshot: Improvement strategies

Table 5 Proportion of primary care physicians who responded that the following strategies are the top priority in order to improve quality of care and patient access, by jurisdiction [d]

Physicians identified patient requests as biggest barrier to reducing low-value or potentially wasteful care in Canada

Provincial and territorial snapshot: Barriers to reducing low-value care

Table 56 Proportion of primary care physicians who responded that the following are major barriers to reducing low-value or potentially wasteful care, by jurisdiction [d]

Barriers	N.L.	P.E.I.	N.S.	N.B.	Que.	Ont.	Man.	Sask.	Alta.	B.C.	Terr.	Can.	CMWF Avg.
Patients expect to receive every test or treatment they want	63		61	58	35	71	53	58	65	57	62	58	57
Lack of time for shared decision making with patients	37		42	38	28	41	41	34	29	42	51	37	41
Medical malpractice environment	25		31		13	38		25		26		27	40
Lack of objective information that helps determine whether patient will benefit from service	24			23	23	23	24	22		28		23	27

Notes

The coefficient of variation (CV) is between and use with caution.

d: Non-directional or no statistical tests were performed to compare with the CMWF average (e.g., territories).

The CMWF average was calculated by adding the results from the 11 countries and dividing by the number of countries. The Canadian average represents the average experience of Canadians (as opposed to the mean of provincial and territorial results).

Methodology notes

The CMWF's 2019 International Health Policy Survey of Primary Care Physicians includes responses from primary care physicians in 11 countries: Australia, Canada, France, Germany, the Netherlands, New Zealand, Norway, Sweden, Switzerland, the United Kingdom and the United States.

More detailed methodology notes, including a complete list of response rates from all countries surveyed, are available online.

In Canada, Social Sciences Research Solutions (SSRS) conducted mail and online surveys from January 29 to June 3, 2019, for the provinces (except P.E.I.), and censuses of P.E.I. and the territories from February 27 to July 30, 2019. In addition to the base sample funded by the CMWF, sample sizes were increased in Quebec and Ontario with funding from provincial organizations, and in the rest of the provinces and territories with funding from CIHI. Primary care physicians were randomly selected in the provinces (except P.E.I.); all primary care physicians in P.E.I. and the territories were invited to participate. To encourage participation, an incentive cheque of $25 or $100 was provided for each primary care physician selected in the provinces and territories, respectively. In total, there were 2,569 respondents in Canada, for an overall response rate of 39.3%.

Due to small sample sizes in Yukon, the Northwest Territories and Nunavut, the results from the territories are reported together (with permission/support).

Weighting of results

7KH VXUYH\ GDWD IRU &DQDGD ZDV ¿UVW ZHLJKWHG E\ DJH DQ
WKH UHVW RI &DQDGD 7KH ZHLJKWV ZHUH VXEVHTXHQWO\ DGM
jurisdiction among Canadian primary care physicians. Benchmarks for physician distribution
ZHUH GHULYHG IURP WKH &0$ 0DVWHU¿OH -DQXDU\ &DQDG

Averages and trends

For this report, the CMWF average was calculated by adding the results from the 11 countries

Demographics of survey respondents (unweighted)



References

1. Statistics Canada. Section 7: Data quality. Accessed October 21, 2019.
2. Canadian Institute for Health Information. *How Canada Compares: Results From The Commonwealth Fund's 2016 International Health Policy Survey of Adults in 11 Countries*. 2017.
3. Canadian Medical Association. *Family Medicine Profile*. 2018.
4. Canadian Institute for Health Information. *How Canada Compares: Results From The Commonwealth Fund 2015 International Health Policy Survey of Primary Care Physicians*. 2016.
5. Canadian Medical Association. *CMA Physician Workforce Survey, 2017*. 2017.
6. Canadian Institute for Health Information. *How Canada Compares: Results From The Commonwealth Fund's 2017 International Health Policy Survey of Seniors*. 2018.
7. Downar J. Resources for educating, training, and mentoring all physicians providing palliative care. *Journal of Palliative Medicine*. 2018.
8. The College of Family Physicians of Canada. *Standards of Accreditation for Residency Programs in Family Medicine*. 2018.
9. Health Canada. *Fourth Interim Report on Medical Assistance in Dying in Canada*. 2019.
10. Canadian Institute for Health Information. *Unnecessary Care in Canada*.

CIHI Ottawa
495 Richmond Road
Suite 600
Ottawa, Ont.
K2A 4H6
613-241-7860

CIHI Toronto
4110 Yonge Street
Suite 300
Toronto, Ont.
M2P 2B7
416-481-2002

CIHI Victoria