DAD

20 Admission for Kidney Donation From Living Donor (D1002-165) 42
21 T40.7 Poisoning by cannabis Without Corresponding Cannabis Poisoning External Cause Code (X42, X62, Y12) (D1002-167) 43
22 High Vaginal Laceration Without Corresponding CCI Repair Code (D1002-168) 44

43 Project 340 — Missing, Invalid or Unknown Value for Date and Time of Acute Thrombolysis Administration When Administration of Acute Thrombolysis Is Y or P (D1618-121) 66
44 Project 340 — Invalid or Unknown Value for Stroke Symptom Onset Date and Time (D1618-123) 67
45 Project 740 — Project Not Completed When an “Applicable Condition” Code Is Recorded (D1619-137) 68
46

Introduction

As part of the Canadian Institute for Health Information’s (CIHI’s) commitment to quality data, the Discharge Abstract Database (DAD) is routinely analyzed for data quality issues during

OYDQ test number	OYDQ test title	Short description	Rationale
D0301-118	Mother's Health Care Number Recorded on In-Province Newborn's Abstract	When available, the provincial/territorial health care number (HCN) assigned to the newborn should be recorded.	High percentages of newborn abstracts with the mother's HCN recorded as HCN diminish the ability to link records of the newborn discharge and any subsequent discharges.
D0402-64	Unknown Admission Time	Admission Time is unknown.	This field is important for episode building.
D0502-65	Unknown Discharge Time	Discharge Time is unknown.	This field is important for episode building.
D0505-175	Discharge Disposition Inconsistent With Institution To	The Discharge Disposition and the CIHI Institution Type assigned to the Institution To number must match.	Accurate data is required for analysis.
D0701-149	(MODIFIED) Potential Alternate Level of Care (ALC) Under-Reporting	Acute inpatients that are likely ALC but ALC patient service is not recorded.	ALC data is well used at every level of the health service system and across acute and continuing care sectors.
D0703-50	Unknown Weight 0.001 Recorded for Newborns and Neonates Less Than 29 Days	Weight is recorded as unknown for newborns and neonates less than 29 days.	Weight impacts the CMG assignment. A high percentage of abstracts with 0001 (Unknown) weight may indicate facility documentation issues.
D1002-32	Incorrect Infant Status of Singleton Within a Multiparous Delivery Episode	The Diagnosis Code of Z38.– on a newborn's abstract indicates the plurality of birth (singleton, twin, triplet, etc.), the same number of newborn abstracts should be linked to the mother's abstract.	Research on multiple birth outcomes is adversely affected by incorrect data.
D1002-52	Post-Procedure Disorder Codes Recorded Without an External Cause Code	All post-procedural disorder codes require an external cause code (Y60–Y84 or V01–X59).	Post-procedural codes are used in reports which are provided to external clients.
D1002-127	Acute Myocardial Infarction Assigned Diagnosis Type (3)	Incorrect Diagnosis Typing for Acute Myocardial Infarction coding	Impacts important Organisation for Economic Co-operation and Development (OECD) indicator.

OYDQ test number	OYDQ test title	Short description	Rationale
D1002-186 **NEW**	Confrmed or Suspected COVID-19 in Obstetrics Delivered	When an obstetrics **delivered** patient is diagnosed with confrmed or suspected COVID-19, O98.5– *viral diseases complicating pregnancy, childbirth and the puerperium* with the sixth digit 1 or 2 must be assigned as a signifcant diagnosis type.	Accurate data is required for in-depth analyses on the impact of COVID-19 on health care systems.
D1002-187 **NEW**	Confrmed or Suspected COVID-19 in Obstetrics Undelivered	When an obstetrics **undelivered** patient is diagnosed with confrmed or suspected COVID-19, O98.5– *Other viral diseases complicating pregnancy, childbirth and the puerperium* with the sixth digit 3 or 4 must be assigned as a signifcant diagnosis type.	Accurate data is required for in-depth analyses on the impact of COVID-19 on health care systems.
D1102-111	Repair High Vaginal Laceration Without a Corresponding Diagnosis Code	Surgical repair of high vaginal laceration without a corresponding Diagnosis Code	Impacts important patient safety indicator.
D1102-180	Cementless Hip/Knee Replacement CCI Codes and the Presence of Cement Sticker	When a cement product number is present, the 10th digit of the corresponding hip/knee replacement CCI code must be “N” or “Q” — with synthetic material.	Accurate data is required for in-depth analyses on hip/knee replacements.
D1103-172	Incorrect Status Attribute With 3.IP.10.VX	Coronary angiography Intervention Code with incorrect Status Attribute	Accurate data is required for in-depth analyses on coronary angiography.
D1103-173	Incorrect Status Attribute With 3.IP.10.VC and/or 3.IP.10.VZ	Intervention Code 3.IP.10. VC and/or 3.IP.10.VZ are recorded AND Status Attribute is DX, I or UN. Status Attribute values DX, I and UN are for use only with 3.IP.10.VX (coronary angiography).	Accurate data is required for in-depth analyses on coronary angiography.
D1104-82	Incorrect Location Attribute With 3.IP.10.VX	Coronary angiography Intervention Code with incorrect Location Attribute	Accurate data is required for in-depth analyses on coronary angiography.

OYDQ test number	OYDQ test title	Short description	Rationale
D1104-174	Incorrect Location Attribute With 3.IP.10.VC and/or 3.IP.10.VZ	Intervention Code 3.IP.10.VC and/or 3.IP.10.VZ is recorded AND Location Attribute is CB, FY, LY, RY or U. Location Attribute values CB, FY, LY, RY and U are for use only with 3.IP.10.VX (coronary angiography).	Accurate data is required for in-depth analyses on coronary angiography.
D1105-86	Extent Attribute UN *(Unknown)* With Hip Replacement	The Extent Attribute UN *(Unknown)* should be used rarely for the implantation of hip prosthesis as the documentation should identify the components implanted.	Attributes are used by CIHI to report on hip replacements.
D1105-166	Incorrect Value of Extent Attribute for Hip Arthroplasty Codes 1.VA.53.^^	When recording hip arthroplasty with implantation prosthetic device codes 1.VA.53.^^, ensure that the value of the extent attribute matches the corresponding Intervention Code.	Generate correct procedure types for primary hip arthroplasty for the Canadian Joint Replacement Registry (CJRR).
D1113-35	Three or More OOH Intervention Episodes in One Day	Recording the Intervention Episode Start Date multiple times for OOH interventions may result in erroneously increasing the number of OOH intervention episodes performed.	Impacts intervention event factor. Intervention count is used in Resource Intensity Weight assignment.
D1515-177	Legal Status Upon Arrival to ED Must Not Be Blank for Non-NACRS L3 ED Facilities	Legal Status Upon Arrival to ED is mandatory for facilities not also reporting NACRS Level 3 ED data.	Accurate data is important for data analysis.
D1515-178	Legal Status Upon Arrival to ED Must Be Blank for NACRS L3 ED Facilities	Legal Status Upon Arrival to ED must be blank for facilities also reporting NACRS Level 3 ED data.	Accurate data is important for data analysis.
D1515-179	Legal Status Upon Arrival to ED Without Entry Code E — Emergency	Legal Status Upon Arrival to ED must be blank if patient was not admitted via the ED.	Accurate data is important for data analysis.
D1618-99	Project 340 — Project Not Completed When an “Applicable Condition” Code Is Recorded (D1618-99)	When a stroke Diagnosis Code is recorded, the Project Number 340 must be completedct	

OYDQ test number	OYDQ test title	Short description	Rationale
D1618-103	Project 340 — Not Applicable, Unknown or Invalid Value for *Antithrombotic Medication* (D1618-103)		

OYDQ test number	OYDQ test title	Short description	Rationale
D1620-155	Project 440 — Invalid Value for CTA, CTP or MRA Scan Performed Prior to Start of Endovascular Thrombectomy Intervention	When Special Project 440 Endovascular Thrombectomy is recorded, it is mandatory to record Field 01 CTA, CTP or MRA Scan Performed Prior to Start of Endovascular Thrombectomy Intervention.	Endovascular thrombectomy is a high-priority health initiative.
D1620-156	Project 440 — Invalid Value for Date and Time of Qualifying Scan Prior to Endovascular Thrombectomy Intervention When CTA, CTP or MRA Scan Performed Prior to Start of Endovascular Thrombectomy Intervention Is P or Y	When Special Project 440 Endovascular Thrombectomy is recorded and Field 01 is recorded with P (Yes, prior) or Y (Yes), fields 02–09 Date and Time of Qualifying Scan Prior to Endovascular Thrombectomy Intervention must be a valid date and time or 99999999 (unknown).	Endovascular thrombectomy is a high-priority health initiative.
D1620-157	Project 440 — Invalid Value for Date and Time of Arterial Puncture	When Special Project 440 Endovascular Thrombectomy is recorded, fields 10–15 Date and Time of Arterial Puncture must be a valid date and time or 999999 (unknown).	Endovascular thrombectomy is a high-priority health initiative.
D1620-158	Project 440 — Invalid Value for First Reperfusion Achieved	When Project 440 Endovascular Thrombectomy is recorded, it is mandatory to record Field 16 First Reperfusion Achieved.	Endovascular thrombectomy is a high-priority health initiative.
D1620-159	Project 440 — Invalid Value for Date and Time of First Reperfusion Achieved When First Reperfusion Achieved Is Y	When Special Project 440 Endovascular Thrombectomy is recorded and Field 16 First Reperfusion Achieved is recorded with Y (Yes), Fields 17–22 Date and Time of First Reperfusion Achieved must be a valid date and time or 999999 (unknown).	Endovascular thrombectomy is a high-priority health initiative.

Open-year data quality tests

1 Potential Extra Abstracts (D0103-18)

Criteria	Description
Selection criteria	Abstracts where the values recorded in the group of data elements below are the same in more than one abstract.
Data elements	Province/Territory, Institution Number, Health Care Number, Birthdate, Gender, Postal Code, Admission Date, Admission Time, Discharge Date, Discharge Time, Diagnosis Code (MRDX), Intervention Code (principal), Weight.

2 Incomplete Linkage of Mothers and Babies (D0112-23)

Rule

The Maternal/Newborn Chart Number on the mother’s record must be the same as the Chart Number recorded on her newborn’s record. The Maternal/Newborn Chart Number on the newborn’s record must be the same as the Chart Number recorded on his or her mother’s record.

Criteria	Description
Selection criteria	Diagnosis Code on mother’s record is Z37.0–, Z37.2–, Z37.3–, Z37.5–, Z37.6– or Z37.9– (delivery)

3 Mother’s Health Care Number Recorded on Out-of-Province Newborn’s Abstract (D0301-117)

Rule

When available, record the provincial/territorial health care number (HCN) assigned to the newborn.

Alberta, Northwest Territories and Yukon:

- When the newborn’s HCN is not available, record the mother’s HCN.
- When the mother’s HCN is not available, record 1 (not applicable) for out-of-province newborns.

New Brunswick:

- When the mother’s HCN is not available for out-of-province newborns, record 1 (not applicable).

Newfoundland and Labrador, P.E.I., Ontario, British Columbia and Nunavut:

- The mother’s health care number cannot be recorded as the health care number on the newborn’s abstracts.
- When the newborn’s HCN is not available, record 1 (not applicable) for out-of-province newborns.

Nova Scotia and Saskatchewan:

- When the HCN for an out-of-province newborn is not available, record 1 (not applicable).

Manitoba:

- When the newborn’s Out of Province HCN is not available, record the mother’s HCN on the newborn’s abstract.

Note: This test will be completed for all out-of-province newborns’ abstracts with a valid HCN. A high percent of out-of-province newborn abstracts with the mother’s HCN recorded as HCN may indicate a need to investigate practices around the capture of out-of-province HCNs for newborns.

Criteria	Description
Selection criteria	**Inclusions:** The abstracts of newborns where • Entry Code is N (Infant born alive within the reporting facility), AND • Province/Territory Issuing HCN is not the same as the province/territory of the reporting facility, AND • HCN has a valid format, AND • Province/Territory Issuing HCN and HCN are equal to the Province/Territory Issuing HCN and HCN on a mother's abstract from the same reporting facility. The abstracts of mothers are used to identify newborns' abstracts with the mother's HCN recorded as HCN. The selection criteria for mothers' abstracts are • One of the Diagnosis Codes is Z37.0–, Z37.2–, Z37.3–, Z37.5–, Z37.6– or Z37.9– (delivery), AND • Most Responsible Diagnosis Code is not O02.– to O05 (abortive outcome), AND • One of the Intervention Codes is 5.MD.50.^^ to 5.MD.60.^^ (delivery). **Exclusions:** 1. Province/Territory Issuing Health Care Number is 99 (not applicable) or CA (Canada). 2. Entry Code is S (Stillborn). 3. Admission Category is R (Cadaveric Donor). 4. HCN is 0 (Insured resident of reporting province/territory but HCN is not available) 5. Or 1 (Not applicable) or HCN has invalid format.
Data elements	Province/Territory, HCN, Province/Territory Issuing HCN, Entry Code, Admission Category, Diagnosis Code, Intervention Code
Correct case example	When available, the provincial/territorial HCN assigned to the newborn is recorded. When the newborn's HCN is not available, record 1 (not applicable) for out-of-province newborn if the province/territory of the reporting facility is Newfoundland and Labrador, P.E.I., Nova Scotia, Ontario, Saskatchewan, British Columbia and Nunavut.
Reference	DAD Abstracting Manual: Group 03 — Patient/Client Demographics, Field 01 — Health Care Number

4 Mother’s Health Care Number Recorded on In-Province Newborn’s Abstract (D0301-118)

Rule

When available, record the provincial/territorial health care number (HCN) assigned to the newborn.

Alberta, Northwest Territories and Yukon:

- When the newborn’s HCN is not available, record the mother’s HCN.
- When the mother’s HCN is not available, record 0 (HCN not available) for provincial/territorial residents.

Newfoundland and Labrador, P.E.I., Ontario, British Columbia and Nunavut:

- The mother’s health care number cannot be recorded as the health care number on the newborn’s abstracts.
- When the newborn’s HCN is not available, record 0 (HCN not available) for provincial/territorial residents.

Nova Scotia, Manitoba and Saskatchewan:

- The newborn’s HCN must always be recorded for provincial/territorial residents.

Note: This test will be completed for all in-province newborns’ abstracts with a valid HCN. A high percent of in-province newborn abstracts with the mother’s HCN recorded as HCN may indicate a need to investigate practices around the capture of HCNs for newborns.

Criteria	Description
Selection criteria	**Inclusions:** The abstracts of newborns where: • Entry Code is N (Infant born alive within the reporting facility), AND • Province/Territory Issuing HCN is the same as the province/territory of the reporting facility, AND • HCN has a valid format, AND • Province/Territory Issuing HCN and HCN are equal to the Province/Territory Issuing HCN and HCN on a mother's abstracts from the same reporting facility. The abstracts of mothers are used to identify newborns' abstracts with the mother's HCN recorded as HCN. The selection criteria for mothers' abstracts are • One of the Diagnosis Codes is Z37.0–, Z37.2–, Z37.3–, Z37.5–, Z37.6– or Z37.9– (delivery), AND • Most Responsible Diagnosis Code is not O02.– to O05 (abortive outcome), AND • One of the Intervention Codes is 5.MD.50.^^ to 5.MD.60.^^ (delivery). **Exclusions:** 1. Province/Territory Issuing Health Care Number is 99 (not applicable) or CA (Canada). 2. Entry Code is S (Stillborn). 3. Admission Category is R (Cadaveric Donor). 4. HCN is 0 (Insured resident of reporting province/territory but HCN is not available) or 1 (Not applicable) or HCN has invalid format.
Data elements	Province/Territory, HCN, Province/Territory Issuing HCN, Entry Code, Admission Category, Diagnosis Code, Intervention Code
Correct case example	When available, the provincial/territorial HCN assigned to the newborn is recorded. When the newborn's HCN is not available, record 0 (HCN not available) for in-province newborn if the province/territory of the reporting facility is Newfoundland and Labrador, P.E.I., Ontario, British Columbia and Nunavut.
ect c)4.(ne74.)J7(e-S)M 1482BCT 1 19 0 0 02.216003417694 286AD EEnuerriTanAct6ntario,Btish Columbia and Nunavt.	

5

6 Unknown Discharge Time (D0502-65)

Rule

Discharge Date/Time is the date and time when the patient was formally discharged.

- Record the date/time the patient was formally discharged and physically left the bed in the nursing unit.
-

7 Discharge Disposition Inconsistent With Institution To (D0505-175)

Rule

The Discharge Disposition and the CIHI Institution Type assigned to the Institution To number must match.

Note: There are known provincial variations. Ontario and Alberta clients should refer to the Notes tab in their OYDQ report.

Criteria	Description
Selection criteria	Abstracts where Discharge Disposition is not consistent with CIHI Institution Type (derived) assigned to Institution To number: Where • Discharge Disposition 04 is not mapped to CIHI Institution Type 0, 6, 8, F, J, M, N, U or Blank OR • Discharge Disposition 10 is not mapped to CIHI Institution Type 1, 2, 3, 5, 9, S or U OR • Discharge Disposition 20 is not mapped to CIHI Institution Type E, A, N, 6 or U OR • Discharge Disposition 30 is not mapped to CIHI Institution Type 4, T, P or U OR • Discharge Disposition 40 is not mapped to CIHI Institution Type G, H or U
Data elements	Discharge Disposition, Institution To
Correct case example	Discharge Disposition 10 — Inpatient Care is recorded along with an Institution To number assigned to CIHI Institution Type 1 — Acute Inpatient Care.
Reference	• Jurisdictional Disposition reference guides can be downloaded via the DAD/NACRS Abstracting Manual link in My Services (with the exception of Yukon and the Northwest Territories). The Instructions tab describes how to use the Disposition Reference Guide while the third tab provides mapping of DAD Discharge Dispositions to province-/territory-specific levels of care. Alberta facilities should refer to the Alberta Supplement to 2018–2019 DAD manual. • DAD Abstracting Manual: Group 05 — Separation, Field 04 Institution To, Field 05 Discharge Disposition • Job Aid — Ensuring Accurate Discharge/Visit Disposition and Institution To Assignment

8 (MODIFIED) Potential Alternate Level of Care (ALC) Under-Reporting (D0701-149)

Rule

When a patient is occupying a bed in a facility and does not require the intensity of resources/services provided in that care setting (acute, chronic or complex continuing care, mental health or rehabilitation), the patient must be designated ALC at that time by the most appropriate care team member, which may be a physician, long term care assessor, patient care manager, discharge planner or other care team member.

The decision to assign ALC status is a clinical responsibility. In order to enter the ALC patient service (99) on the DAD abstract, there must be clear, consistent documentation by the clinical staf, preferably on an approved ALC Designation form.

Note: Accurate ALC reporting is key to monitoring and improving access to services, patient fow and outcomes in acute care. This test estimates the percent of acute inpatients that are likely ALC but have no ALC days recorded.

Criteria	Description
Selection criteria	Acute inpatients that are likely ALC but have no ALC days recorded: • Calculated Length of Stay (LOS) is greater than Expected Length of Stay (ELOS)*, adjusted for Case Mix Groups (CMG); and • Discharge Disposition is 20 (ED and Ambulatory Care), 30 (Residential Care), 40 (Group/Supportive Living), 04 (Home with Support/Referral) or 90 (Correctional Facility); and

9 Unknown Weight 0.001 Recorded for Newborns and Neonates Less Than 29 Days (D0703-50)

Rule

The weight of a newborn or neonate less than 29 days on admission to the facility must be recorded.

Every effort should be made to record the admission weight since Weight is required for Case Mix Group (CMG+) assignment.

Criteria	Description
Selection criteria	The abstracts of newborns and neonates where Age Code is D (Days) or B (Newborn or Stillbirth) AND Age Unit is 0–28 AND Weight is 0.001 AND Entry Code is not equal to S (Stillborn)
Data elements	Entry Code, Age (Derived), Weight

10 Incorrect Infant Status of Singleton Within a Multiparous Delivery Episode (D1002-32)

Rule

According to the Canadian Coding Standards, every newborn record must include a code from Z38.– *Liveborn infants according to place of birth* to indicate the plurality of birth:

- A liveborn singleton is assigned a code from Z38.0– to Z38.2–.
- Liveborn twins, triplets or other multiple births are assigned a code from Z38.3– to Z38.8–.

A multiple birth newborn record must not have a code from Z38.0– to Z38.2– (singleton) recorded.

Most multiple births are delivered on the same date; however, some multiple births can occur on different dates. The codes Z38.3– to Z38.8– describe the plurality of the pregnancy and apply even when the births occur on different days or at different locations and/or when 1 or more of the babies are stillborn.

Note: This analysis focuses on multiple births delivered on the same date. Clients may also perform analyses on different delivery dates, different delivery locations and where 1 or more newborns are stillborn.

Criteria	Description
Selection criteria	Newborn abstracts with Diagnosis Code Z38.0– to Z38.2– (exclude P96.4 *Termination* as Most Responsible Diagnosis) AND More than one Maternal/Newborn Chart Number recorded with the same admission date.
Data elements	Entry Code, Chart Number, Maternal/Newborn Chart Number, Admission Date, Diagnosis Code
Correct case examples	Mother: Chart Number: 8866766 Baby A: Admission Date: 2013/11/01 Entry Code: N Chart Number: 123455 Maternal/Newborn Chart Number: 8866766 Z38.300 *Twin, born in hospital, delivered vaginally, product of both spontaneous (NOS) ovulation and conception*.

Criteria	Description
Correct case examples (continued)	Baby B: Admission Date: 2013/11/01 Entry Code: N Chart Number: 123456 Maternal/Newborn Chart Number: 8866766 Z38.310 *Twin, born in hospital, delivered by caesarean, product of both spontaneous (NOS) ovulation and conception*
Reference	Canadian Coding Standards: Diagnosis Typing Definitions for DAD

11 Post-Procedural Disorder Codes Recorded Without an External Cause Code (D1002-52)

Rule

All post-procedural disorder codes (see Appendix A) require an external cause code (Y60–Y84 or V01–X59).

Criteria	Description
Selection criteria	Abstracts with a post-procedural disorder Diagnosis Code (see Appendix A) without an External Cause Code (Y60–Y84 or V01–X59).
Data element	Diagnosis Code
Correct case example	K91.42 (M) *Malfunction of colostomy stoma, not elsewhere classified* **[Diagnosis Cluster A]** Y83.3 (9) *Surgical operation with formation of external stoma as the cause of abnormal reaction of the patient, or of later complication, without mention of misadventure at the time of the procedure* **[Diagnosis Cluster A]**
References	Canadian Coding Standards: Post-Intervention Conditions Self-Learning Product: Classifying Post-Intervention Conditions: ICD-10-CA Code Assignment

12 Acute Myocardial Infarction Assigned Diagnosis Type (3) (D1002-127)

Rule

A myocardial infarction within the acute phase is always assigned a significant Diagnosis Type (M, 1, 2, W, X, Y) per the coding standard Acute Coronary Syndrome (ACS).

Exceptions:

1. When a patient is readmitted with a diagnosis classifiable to category I22.–

   Subsequent myocardial infarction, a code from category I21.– *Acute myocardial infarction* may be assigned as an optional Diagnosis Type (3)/other problem to indicate the site of the original MI.
2. When I21.– is recorded with a Diagnosis Prefix Q, Diagnosis Type (3) may be appropriate for the case.
3. Obstetric cases where O99.4– is recorded, I21.– is assigned a Diagnosis Type (3) to identify MI as the specific condition.

Criteria	Description
Selection criteria	Abstracts where Diagnosis Code I21 is recorded as Diagnosis Type (3) AND Diagnosis Prefix is not Q AND Diagnosis Code I22.– AND O99.4– is not recorded on the same abstract.
Data elements	Diagnosis Code, Diagnosis Type, Diagnosis Prefix
Correct case examples	I21.1 (M) Acute transmural myocardial infarction of inferior wall R94.30 (3) Electrocardiogram suggestive of ST segment elevation myocardial infarction [STEMI]
Reference	Canadian Coding Standards: Acute Coronary Syndrome

13 MRSA and VRE Infections — Missing Additional Code for Site of Infection (D1002-134)

Rule

When a current infection is documented as due to one of the “super bugs” referred to as MRSA (methicillin-resistant) or VRE (vancomycin-resistant enterococcus), it is mandatory to assign the codes that identify the site of the infection, the causative organism and the drug resistance (as described below). The Diagnosis Cluster is mandatory to link all the codes together.

MRSA infection:

- A code that identifes the site of the infection;
- B95.6 *Staphylococcus aureus as the cause of diseases classifed to other chapters*; and
- U82.1 *Resistance to methicillin*

VRE infection:

- A code that identifes the site of the infection;
- B95.21 *Enterococcus as the cause of diseases classifed to other chapters*; and
- U83.0 *Resistance to vancomycin*

Criteria	Description
Selection criteria	Abstracts where Diagnosis Codes U82.1 *Resistance to methicillin* AND B95.6 *Staphylococcus aureus as the cause of diseases classifed to other chapters* are recorded with the same Diagnosis Cluster value AND there is no other code in the same Diagnosis Cluster OR Abstracts where Diagnosis Codes U83.0 *Resistance to vancomycin* and B95.21 *Enterococcus as the cause of diseases classifed to other chapters* are recorded with the same Diagnosis Cluster value AND There is no other code in the same Diagnosis Cluster.
Data elements	Diagnosis Code, Diagnosis Cluster
Correct case examples	M00.01 (M) *Staphylococcal arthritis and polyarthritis, shoulder region* **[Diagnosis Cluster A]** AND B95.6 (3) *Staphylococcus aureus as the cause of diseases classifed to other chapters* **[Diagnosis Cluster A]** AND U82.1 (1) *Resistance to methicillin* **[Diagnosis Cluster A]**
References	Canadian Coding Standards: Drug-Resistant Microorganisms Diagnosis Cluster

14 Incorrect Coding of Post-Intervention Sepsis (D1002-136)

Rule

nec ndi i n, Wh pri ary Ա ֏ ֏

- T84.64 Infection and inflammatory reaction due to internal fixation device of tibia and fibula
- T84.65 Infection and inflammatory reaction due to internal fixation device of bones of foot
- T84.68 Infection and inflammatory reaction due to internal fixation device of bones at other site
- T84.69 Infection and inflammatory reaction due to internal fixation device of bones of limb NOS
- T84.7 Infection and inflammatory reaction due to other internal orthopaedic prosthetic devices, implants and grafts
- T85.7 Infection and inflammatory reaction due to other internal prosthetic devices, implants and grafts

Criteria	Description
Selection criteria	Abstracts where any one of the following Diagnosis Codes A02.1, A03.9, A20.7, A21.7, A22.7, A23–, A24.1, A26.7, A28.0, A28.2, A32.7, A39.2, A39.3, A39.4, A40.–, A41.–, A42.7, A54.86, B37.7 is recorded with the same Diagnosis Cluster value as Y83–Y84 AND There is no T-code (from the list included in the Rule above) recorded within the same Diagnosis Cluster.
Data elements	Diagnosis Code, Diagnosis Cluster
Correct case examples	T81.4 (2) *Infection following a procedure, not elsewhere classified* **[Dx Cluster A]** A41.9 (3) *Sepsis, unspecified* **[Dx Cluster A]** Y83.8 (9) *Other surgical procedures as the cause of abnormal reaction of the patient, or of later complication, without mention of misadventure at the time of the procedure* **[Dx Cluster A]**
References	Canadian Coding Standards: Septicemia/Sepsis Post-Intervention Conditions Assignment of Additional Codes for Specificity Appendix E — Tips for Coders: Selecting the Primary Code for a Post-Intervention Condition

15 More Than One Outcome of Delivery (Z37) Code (D1002-148)

Rule

When a delivery occurs during an episode of care, it is mandatory to assign **one Diagnosis** Code from category Z37 *Outcome of delivery* on the mother's abstract. The outcome of delivery codes are broken down into subcategories by number of births (singleton versus multiple) and whether the outcome is a livebirth or stillbirth. Only **one** outcome of delivery code is assigned on the mother's abstract and is dependent on the circumstances of the specifc case.

Criteria	Description
Selection criteria	Abstracts with more than one Z37.– Diagnosis Code.
Data element	Diagnosis Code
Correct case examples	O36.491 *Maternal care for intrauterine death, unspecifed trimester, delivered, with or without mention of antepartum condition* O30.001 *Twin pregnancy, delivered, with or without mention of antepartum condition* Z37.300 *Twins, one liveborn and one stillborn, pregnancy resulting from both spontaneous ovulation and conception*
Reference	ICD-10-CA coding convention and rule

16 Neurologically Determined Death Not Assigned Diagnosis Type (3) (D1002-152)

Rule

The Diagnosis Code G93.81 *Neurologically determined death* must be assigned as a Diagnosis Type (3), mandatory, when there is documentation of brain death by a designated physician.

Criteria	Description
Selection criteria	Abstracts where Diagnosis Code G93.81 is not assigned as a Diagnosis Type (3) **Exclusion:** Diagnosis Type (0)
Data elements	Diagnosis Code, Diagnosis Type
Correct case example	G93.81 (3) *Neurologically determined death* assigned
Reference	Canadian Coding Standards: Neurologically Determined Death

Criteria	Description
Selection criteria	Abstracts with a Diagnosis Code T40.0, T40.1, T40.2–, T40.3, T40.4– or T40.6 **without** external cause code X42, X62 or Y12 **Exclusions:** T40.0, T40.1, T40.2–, T40.3, T40.4– or T40.6 **with** X85 *Assault by drugs, medicaments and biological substances*
Data elements	Diagnosis Code
Correct case examples	T40.40 (M) Poisoning by fentanyl and derivatives X42 (9) Accidental poisoning by and exposure to narcotics and psychodysleptics [hallucinogens], not elsewhere classified
References	ICD-10-CA Table of Drugs Coding Standard: *Adverse Reactions in Therapeutic Use Versus Poisoning* Coding Standards Appendix A: Opioid overdose Classifications bulletin: Opioid Overdose Coding Direction

18 Alcohol Poisoning External Cause Code Without Corresponding Alcohol Poisoning T-Code (D1002-163)

Rule

When one of the following alcohol poisoning external cause codes is recorded:

- X45 Accidental poisoning by and exposure to alcohol
- X65 Intentional self-poisoning by and exposure to alcohol
- Y15 Poisoning by and exposure to alcohol, undetermined intent

The corresponding alcohol poisoning T-code from category T51.– *Toxic effect of alcohol* must also be recorded.

Criteria	Description
Selection criteria	Abstracts with external cause code X45, X65 or Y15 **without** a Diagnosis Code from category T51.–
Data element	Diagnosis Code
Correct case examples	T51.9 (M) Toxic effect of alcohol, unspecified X45 (9) Accidental poisoning by and exposure to alcohol
References	ICD-10-CA Table of Drugs Coding Standard: *Adverse Reactions in Therapeutic Use Versus Poisoning*

19 Alcohol Poisoning T-Code Without Corresponding Alcohol Poisoning External Cause Code (D1002-164)

Rule

When a code from category T51.– *Toxic effect of alcohol* is recorded, one of the following poisoning external cause codes must also be recorded:

•

20 Admission for Kidney Donation From Living Donor (D1002-165)

Rule

When a patient is admitted for the sole purpose of donating a kidney, the most responsible diagnosis (MRDX) should be Z52.4 *Kidney donor* and the Intervention Code must be 1.PC.58.^^-XX-J *Procurement, kidney,* by approach, *from living donor*.

CCI provides an Excludes instruction at 1.PC.89.^^ *Excision total, kidney* to alert coders that procurement of kidney from a living donor is classified to 1.PC.58.^^ *Procurement, kidney*.

Criteria	Description
Selection criteria	Abstracts where MRDx is Diagnosis Code Z52.4 and Intervention Code is 1.PC.89.^^ or 1.PC.91.^^
Data elements	Diagnosis Code, Diagnosis Type, Intervention Code
Correct case examples	Z52.4 (M) Kidney donor 1.PC.58.LB-XX-J Procurement, kidney, open abdominal approach, from living donor
References	ICD-10-CA CCI

21 T40.7 *Poisoning by cannabis* Without Corresponding Cannabis Poisoning External Cause Code (X42, X62, Y12) (D1002-167)

Rule

When the Diagnosis Code T40.7 *Poisoning by cannabis (derivatives)* is recorded, one of the following cannabis poisoning external cause codes must also be recorded:

- X42 *Accidental poisoning by and exposure to narcotics and psychodysleptics [hallucinogens], not elsewhere classifed;* or
- X62 *Intentional self-poisoning by and exposure to narcotics and psychodysleptics [hallucinogens], not elsewhere classifed;* or
- Y12 *Poisoning by and exposure to narcotics and psychodysleptics [hallucinogens], not elsewhere classifed, undetermined intent.*

When the chart review reveals that

1. The case documentation meets the definition of cannabis poisoning, then correct the case by assigning the applicable cannabis poisoning external cause code X42, X62 or Y12 (code could be missing or could be the wrong poisoning external cause code); or
2. The case documentation did not meet the criteria to classify to T40.7 *cannabis (derivatives)*, then remove the T-code representing cannabis poisoning. For example, a diagnosis of nausea due to correct use of prescribed medical cannabis is **not** classifed to T40.7 and Y49.6 *Psychodysleptics [hallucinogens] causing adverse effects in* . In such a circumstance, per the Canadian Coding Standards, a code for the specifc adverse reaction(s) is assigned (not T40.7) followed by Y49.6.

Criteria	Description
Selection criteria	Acute care abstracts with a Diagnosis Code T40.7 without external cause code X42, X62 or Y12 **Exclusions:** T40.7 with X85 *Assault by drugs, medicaments and biological substances*
Data element	Diagnosis Code
Correct case example	T40.7 (M) Poisoning by cannabis (derivatives) X42 (9) Accidental poisoning by and exposure to narcotics and psychodysleptics [hallucinogens], not elsewhere classifed
References	ICD-10-CA Table of Drugs Coding Standard: *Adverse Reactions in Therapeutic Use Versus Poisonings*

22 High Vaginal Laceration Without Corresponding CCI Repair Code (D1002-168)

Rule

For obstetric delivered or obstetric postpartum episodes of care, the Diagnosis Code O71.401 or O71.404 is expected to also have the corresponding Intervention Code 5.PC.80.JU *Surgical repair, postpartum, of current obstetric high vaginal laceration* recorded.

Notes

- When chart review reveals a high vaginal laceration repair was performed and the corresponding CCI repair code is missing or the wrong CCI repair code was assigned, correct the case by assigning 5.PC.80.JU and, if applicable, removing the wrong CCI repair code.
- When chart review reveals there was no documentation of a high vaginal laceration and,

23

24 I64 *Stroke, not specified as haemorrhage or infarction* Recorded (D1002-170)

Rule

Classify a diagnosis of “stroke” or “cerebrovascular accident” as

- A **hemorrhagic stroke** when there is documentation to support there was subarachnoid hemorrhage (I60.–) or intracerebral hemorrhage (I61.–); or
- An **ischemic stroke** (I63.–) when a hemorrhagic stroke has been ruled out.

Notes

I64

26 U07.4 *Post COVID-19 condition* Assigned Diagnosis Type M, 1, 2, W, X, Y (D1004-185)

Rule

When U07.4 *Post COVID-19 condition* is assigned on an abstract, the diagnosis type must be (3).

Note: U07.4 *Post COVID-19 condition* is to be used in multiple coding to identify a case in which the physician/primary care provider documented that a condition or symptom has a relationship or association with past COVID-19.

Criteria	Description
Selection criteria	Abstracts where U07.4 is assigned diagnosis type M, 1, 2, W, X, Y
Data elements	Diagnosis Type, Diagnosis Code
Correct case example	U07.4 is assigned diagnosis type (3).
Reference	Bulletin November 12, 2020: ICD-10-CA coding direction for post COVID-19 condition and personal history of COVID-19

27 Con

28 Confirmed or Suspected COVID-19 in Obstetrics Undelivered

Rule

When an obstetrics **undelivered** patient is diagnosed with confirmed or suspected COVID-19, O98.5– *Other viral diseases complicating pregnancy, childbirth and the puerperium*

29 Repair High Vaginal Laceration Without a Corresponding Diagnosis Code (D1102-111)

Rule

For obstetrics delivered or obstetric postpartum episodes of care, the Intervention Code 5.PC.80.JU *Surgical repair, postpartum, of current obstetric high vaginal laceration* must have a corresponding Diagnosis Code of O71.401 or O71.404 recorded on the same abstract.

The error can be due to

- An Intervention Code assigned correctly but the Diagnosis Code is missing, OR
- An Intervention Code assigned incorrectly

Criteria	Description
Selection criteria	Intervention Code 5.PC.80.JU *Surgical repair, postpartum, of current obstetric high vaginal laceration* is recorded **without** Diagnosis Code O71.401 or O71.404 *high vaginal laceration* Exclude abortion abstracts with a Diagnosis Code from O08.6–
Data elements	Intervention Code, Diagnosis Code
Correct case examples	5.PC.80.JU *Surgical repair, postpartum, of current obstetric high vaginal laceration* AND O71.401 *Obstetric high vaginal laceration, delivered, with or without mention of antepartum condition*

31 Incorrect Status Attribute With 3.IP.10.VX (D1103-172)

Rule

Status Attribute 0 (Not applicable) must not be assigned with 3.IP.10.VX *coronary arteries, of left heart structures using percutaneous transluminal arterial (retrograde)* . When coronary angiography is performed, the Status Attribute must identify whether the coronary angiography was

- Diagnostic [DX]
- Intraoperative [I]
- Unknown [whether diagnostic or intraoperative] [UN]

Note: The Status Attribute 0 (Not applicable) [i.e., ventriculogram] is for use only with 3.IP.10.VC *transluminal venous (antegrade) approach* and/or 3.IP.10.VZ *arteries of left heart structures using percutaneous transluminal transseptal approach.*

32 Incorrect Status Attribute With 3.IP.10.VC and/or 3.IP.10.VZ (D1103-173)

Rule

The following status attributes **must not** be assigned with 3.IP.10.VC *coronary arteries of right heart structures using percutaneous transluminal venous (antegrade) approach* and/or 3.IP.10.VZ *structures using percutaneous transluminal transseptal approach:*

- Diagnostic [DX]
- Intraoperative [I]
- Unknown [whether diagnostic or intraoperative] [UN]

Note: Status Attribute values DX, I and UN are for use only with 3.IP.10.VX (coronary angiography).

Criteria	Description
Selection criteria	Abstracts where Intervention Code 3.IP.10.VC and/or 3.IP.10.VZ is recorded AND Status Attribute is DX, I or UN.
Data elements	Intervention Code, Location Attribute
Correct case example	3.IP.10.VC *percutaneous transluminal venous (antegrade) approach* Status Attribute: 0 (Not applicable)
Reference	CCI Status Attribute pop-up note

33 Incorrect Location Attribute With 3.IP.10.VX (D1104-82)

Rule

Location Attribute 0 (Not applicable) must not be assigned with 3.IP.10.VX *coronary arteries, of left heart structures using percutaneous transluminal arterial (retrograde)* . Coronary angiography is always performed via an arterial access and as such the Location Attribute must represent arterial access:

- Combined [for chronic total occlusion] [CB]
- Femoral artery (left) (right) [FY]
- Left arm artery [e.g., radial, brachial] [LY]
- Right arm artery [e.g., radial, brachial] [RY]
- Unknown artery [U]

Note: The Location Attribute 0 (not applicable) [i.e., venous approach] is for use only with 3.IP.10.VC *transluminal venous (antegrade) approach* and/or 3.IP.10.VZ *arteries of left heart structures using percutaneous transluminal transseptal approach*.

Criteria	Description
Selection criteria	Abstracts where Intervention Code 3.IP.10.VX is recorded AND Location Attribute is 0.
Data elements	Intervention Code, Location Attribute
Correct case example	3.IP.10.VX *percutaneous transluminal arterial (retrograde) approach* Location Attribute: FY (Femoral artery (left) (right))
Reference	CCI Location Attribute pop-up note

34 Incorrect Location Attribute With 3.IP.10.VC and/or 3.IP.10.VZ (D1104-174)

Rule

The following location attributes representing arterial access **must not** be assigned with 3.IP.10.VC

35 Extent Attribute UN (Unknown) With Hip Replacement (D1105-86)

Rule

Select the Extent Attribute UN (Unknown) only when there is no information to select a VSHFLILF YDOXH IURP WKH H[WHQW DWWULEXWH RSWLRQV 81 8QNQ documentation will identify the component used with a hip replacement procedure 1.VA.53.^^ ,PSODQWDWLRQ RI LQWHUQDO GHYLFH KLS MRLQW

Criteria	Description
Selection criteria	Abstracts where the Intervention Code from rubric 1.VA.53.^^ is recorded with the Extent Attribute UN Exclude hip replacement procedures where Out-of-Hospital Indicator is Y (Yes) or Status Attribute is A (Abandoned)
Data elements	Intervention Code, Extent Attribute, Status Attribute, Out-of-Hospital Indicator
Correct case examples	1.VA.53.LA-PN-N ,PSODQWDWLRQ RI LQWHUQDO GHYLFH KLS MRLQW GHYLFH >IHPRUDO ZLWK DFHWDEXODU@ XVLQJ V\QWKHWLF PDW '\QDJUDIW 2VWHRVHW 6WDWXV $WWULEXWH 5 UHYLVLRQ /RFDWLRQ / OHIW Extent Attribute: FH 0RGXODU EDOO ZLWK RU ZLWKRXW PRGXODU QHF LQWLX >WKLV FRPSRQHQW YDOXH UHTXLUHV 6WDWXV YDOXH 5

36 Incorrect Value of Extent Attribute for Hip Arthroplasty Codes 1.VA.53.^^ (D1105-166)

Rule

When recording hip arthroplasty with implantation prosthetic device codes 1.VA.53.^^, ensure that the value of the extent attribute matches the corresponding Intervention Code.

- For a hip arthroplasty

37 Three or More OOH Intervention Episodes in One Day (D1113-35)

Rule

Interventions are captured in intervention episodes. An intervention episode represents a patient's visit to a physical location where 1 or more interventions may take place. When more than one CCI code is required to capture the interventions performed in a single intervention episode, the Intervention Episode Start Date will be recorded once on the first line of the abstract. Every time an Intervention Episode Start Date is recorded on the abstract, a new intervention episode is derived.

The Out-of-Hospital (OOH) Indicator field indicates that an intervention episode was performed in the day surgery or other ambulatory care setting of another facility during the current inpatient stay in the reporting facility.

38 Legal Status Upon Arrival to ED Must Not Be Blank f

—

39 Legal Status Upon Arrival to ED Must Be Blank for NACRS L3 ED Facilities (D1515-178)

Rule

Legal Status Upon Arrival to ED **must be blank** for facilities also reporting NACRS Level 3 ED data if the patient was admitted via the ED. An entry in this field represents unnecessary data collection and increased coder burden. Data can be obtained via linkage to the corresponding NACRS record.

Criteria	Description
Selection criteria	Abstracts from acute inpatient institutions that report NACRS Level 3 ED data: Where • Entry Code is E (Emergency) AND • Main Patient Service or Service Transfer Service is 64 (Psychiatry) or 65 (Pediatric Psychiatry) AND • Legal Status Upon Arrival to ED is not blank
Data elements	Entry Code, Main Patient Service, Service Transfer Service, Legal Status Upon Arrival to ED
Correct case example	Entry Code E (Emergency) and Main Patient Service 64 (Psychiatry) were recorded on an acute inpatient abstract. The submitting institution does submit NACRS Level 3 ED data. Legal Status Upon Arrival to ED data element remains blank.
Reference	• DAD Abstracting Manual: Group 04 - Admission, Field 06 Entry Code, Group 07 — Patient Service, Field 01 Main Patient Service, Group 08 — Service Transfer, Field 01 Service Transfer Service, Group 15 — Mental Health Indicators, Field 15 Legal Status Upon Arrival to ED • Job Aid — Frequency of Restraint Use and Legal Status Upon Arrival to ED in the DAD

40 Legal Status Upon Arrival to ED Without Entry Code E — Emergency (D1515-179)

Rule

Legal Status Upon Arrival to ED **must be blank** if the patient was not admitted via the ED.

Criteria	Description
Selection criteria	Abstracts from acute inpatient: Where • Entry Code is not E (Emergency) AND • Main Patient Service or Service Transfer Service is 64 (Psychiatry) or 65 (Pediatric Psychiatry) AND • Legal Status Upon Arrival to ED is not blank
Data elements	Entry Code, Main Patient Service, Service Transfer Service, Legal Status Upon Arrival to ED
Correct case example	Main Patient Service 64 (Psychiatry) is recorded on an acute inpatient abstract. The Entry Code is D (Direct). The Legal Status Upon Arrival to ED data element remains blank.
Reference	• DAD Abstracting Manual: Group 04 Admission, Field 06 Entry Code, Group 07 — Patient Service, Field 01 Main Patient Service, Group 08 — Service Transfer, Field 01 Service Transfer Service, Group 15 — Mental Health Indicators, Field 15 Legal Status Upon Arrival to ED • Job Aid

41 Project 340 — Project Not Completed When an “Applicable Condition” Code Is Recorded (D1618-99)

Rule

Special Project 340 Canadian Stroke Strategy Performance Improvement is expected to be completed for all patients who have been diagnosed with an acute/current stroke, and certain other conditions that from an ICD-10-CA classifcation perspective are not classifed as a hemorrhagic, ischemic or unspecifed stroke. The other conditions included in this project are: transient ischemic attack (TIA), transient retinal artery occlusion, intracranial and intraspinal phlebitis and thrombophlebitis, nonpyogenic thrombosis of intracranial venous system and central retinal artery occlusion.

Note: The term **“applicable condition”** is used throughout the stroke projects documentation to refer to the ICD-10-CA codes/conditions included.

Inclusion criteria ICD-10-CA code list:

-

43 Project 340 — Missing, Invalid or Unknown Value for Date and Time of Acute Thrombolysis Administration When Administration of Acute Thrombolysis Is Y or P (D1618-121)

Rule

When Special Project 340 Special Project Canadian Stroke Strategy Performance Improvement is completed, it is mandatory to record Fields 04 to 11 Date and Time of Acute Thrombolysis Administration when Field 03 Administration of Acute Thrombolysis is Y (Yes) or P (Yes, Prior). This feld captures the specifc date and time that a patient received acute thrombolysis. The start date/time for administration of the medication should be recorded in these felds. The year is not being recorded.

Note: A high percent of abstracts with missing (blank), invalid or 99 (unknown) date and time may indicate a need to investigate documentation practices.

Criteria	Description
Selection criteria	An “applicable condition” is recorded AND Field 03 Administration of Acute Thrombolysis is Y (yes) or P (yes, prior) AND 1 or more of the following felds are blank, unknown or invalid: • Fields 04–05 (Month): is blank, is 99 (unknown) or is not 01–12 • Fields 06–07 (Day): is blank, is 99 (unknown) or is not 01–31 • Fields 08–09 (Hour): is blank, is 99 (unknown) or is not 00–23 • Fields 10–11 (Minutes): is blank, is 99 (unknown) or is not 00–59 **Exclusions:** • When acute thrombolysis was given for a condition other than an “applicable condition” • Hemorrhagic strokes (I60 or I61)
Data elements	Project Number, Fields 03 to 11
Reference	Group 16 — Special Projects documentation is now accessible through the DAD/NACRS Abstracting Manual application on CIHI’s website.

44 Project 340 — Invalid or Unknown Valuefor

45 Project 740 — Project Not Completed When an “Applicable Condition” Code Is Recorded (D1619-137)

Rule

Special Project 740 AlphaFIM® is mandatory to report for all DAD acute inpatient admissions of patients 18 years and older with an acute/current stroke, and certain other conditions that from an ICD-10-CA classifcation perspective are not classifed as a hemorrhagic, ischemic or unspecifed stroke. The other conditions included in this project are transient ischemic attack (TIA), intracranial and intraspinal phlebitis and thrombophlebitis and nonpyogenic thrombosis of intracranial venous system.

Note: The term **“applicable condition”** is used throughout the stroke projects documentation to refer to the ICD-10-CA codes/conditions included in in the diferent projects.

Inclusion criteria ICD-10-CA code list:

- I60.– *Subarachnoid haemorrhage*
- I61.– *Intracerebral haemorrhage*
- I63.–
- I64.– *Stroke, not specifed as haemorrhage or infarction*
- I67.6 *Nonpyogenic thrombosis of intracranial venous system*
- G08 *Intracranial and intraspinal phlebitis and thrombophlebitis*
- G45.– *Transient cerebral ischaemic attacks and related syndromes* (**excluding** G45.4 *Transient global amnesia*)
- O22.5– *Cerebral venous thrombosis in pregnancy*
- O87.3– *Cerebral venous thrombosis in puerperium*

Notes:

- Special Project 740 is not completed for the codes H34.0– (another valid type of TIA) and H34.1– .
- There may be cases fagged with this test that do not require completion of Project 740.

Criteria	Description
Selection criteria	Acute care abstracts from Ontario and from the following Newfoundland and Labrador facilities — Health Sciences Centre 00215, St. Clare's Mercy Hospital 00214, Dr. G.B. Cross Memorial Hospital 00064, Burin Peninsula Health Care Centre 00065, Carbonear General Hospital 00060, Central Newfoundland Regional Health Centre 00038, James Paton Memorial Hospital 00049 and Western Memorial Regional Hospital 00004 — where Project 740 is not completed when a Diagnosis Code for an "applicable condition" (see code list above) is recorded and the patient is 18 years or older **Exclusion:** Abstracts where Z51.5– *Palliative care* is assigned
Data elements	Diagnosis Code, Project Number
Reference	Group 16 — Special Projects documentation is now accessible through the DAD/NACRS Abstracting Manual application on CIHI's website.

46 Project 740 — Invalid Value for Documentation of AlphaFIM® Scores (D1619-138)

Rule

When Project 740 AlphaFIM® is completed, it is mandatory to record Field 01 Documentation of AlphaFIM® Scores.

This field captures whether AlphaFIM® scores Projected FIM® — 13 Raw Motor Rating and/or the Projected FIM® — 5 Raw Cognitive Rating were documented for patients with an “applicable condition.”

Criteria	Description
Selection criteria	An “applicable condition” is recorded AND Field 01 Documentation of AlphaFIM® Scores is not equal to Y or N
Data elements	Project Number, Field 01
Reference	Group 16 — Special Projects documentation is now accessible through the

47 Project 740 — Invalid Value for AlphaFIM® Completion Date When Documentation of AlphaFIM® Scores Is Y (D1619-139)

Rule

When Special Project 740 AlphaFIM® is completed and Field 01 Documentation of AlphaFIM® is Y, Fields 02–09 AlphaFIM® Completion Date must be a valid date or 99999999 (unknown).

This field captures the first date when AlphaFIM® scores Projected FIM® — 13 Raw Motor Rating and/or Projected FIM® — 5 Raw Cognitive Rating were documented.

Criteria	Description
Selection criteria	An "applicable condition" is recorded AND Field 01 Documentation of AlphaFIM® Scores is Y AND 1 or more of the following fields are invalid: • Fields 02–05 (Year): not a valid calendar year or not 9999 (unknown) • Fields 06–07 (Month): not a valid calendar month or not 99 (unknown) • Fields 08–09 (Day): not a valid calendar day or not 99 (unknown)
Data elements	Project Number, Field 01, Fields 02–09
Reference	Group 16 — Special Projects documentation is now accessible through the DAD/NACRS Abstracting Manual application on CIHI's website.

48 Project 740 — AlphaFIM® Completion Date Not Between Admission Date and

49 Project 740 — Invalid Value for Projected FIM® — 13 Raw Motor Rating (D1619-142)

Rule

When Special Project 740 AlphaFIM® is completed and Field 01 Documentation of AlphaFIM® is Y, Fields 10–11 Projected FIM® — 13 Raw Motor Rating must be greater than or equal to 13 and less than or equal to 81 or equal to 99 (unknown).

This data element captures the total score documented for a patient's motor functional status from among the relevant tasks of eating, grooming, transfers, locomotion and bowel management.

Criteria	Description
Selection criteria	An "applicable condition" is recorded AND Field 01 Documentation of AlphaFIM® Scores is equal to Y AND Fields 10–11 Projected FIM® — 13 Raw Motor Rating is not greater than or equal to 13 and less than or equal to 81 or equal to unknown value 99
Data elements	Project Number, Field 01, Fields 10–11
Reference	Group 16 — Special Projects documentation is now accessible through the DAD/NACRS Abstracting Manual application on CIHI's website.

51 Project 440 — Invalid Value for CTA, CTP or MRA Scan Performed Prior to Start of Endovascular Thrombectomy Intervention (D1620-155)

Rule

When Special Project 440 Endovascular Thrombectomy is recorded, it is mandatory to record Field 01 CTA, CTP or MRA Scan Performed Prior to Start of Endovascular Thrombectomy Intervention.

This data element captures where the last (most recent) computed tomography angiography (CTA), computed tomography perfusion (CTP) or magnetic resonance angiogram (MRA) scan

52 Project 440 — Invalid Value for Date and Time of Qualifying Scan Prior to Endovascular Thrombectomy Intervention When CTA, CTP or MRA Scan Performed Prior to Start of Endovascular Thrombectomy Intervention Is P or Y (D1620-156)

Rule

When Special Project 440 Endovascular Thrombectomy is recorded and Field 01 CTA, CTP or MRA Scan Performed Prior to Start of Endovascular Thrombectomy Intervention is recorded with P (Yes, prior) or Y (Yes), Fields 02–09 Date and Time of Qualifying Scan Prior to Endovascular Thrombectomy Intervention must be a valid date and time or 99999999 (unknown).

This data element captures the date and time of the last CTA, CTP or MRA scan used to decide whether to attempt an endovascular thrombectomy. This is the date/time of the first slice image, which is electronically stamped on the images when the CTA, CTP or MRA scan is started.

Criteria	Description
Selection criteria	Project Number 440 is recorded And Field 01 CTA, CTP or MRA Scan Performed Prior to Start of Endovascular Thrombectomy Intervention is P (Yes, prior) or Y (Yes) And 1 or more of the following fields are blank or invalid: • Fields 02–03 (Month): not 01–12 or not 99 (unknown) • Fields 04–05 (Day): not 01–31 or not 99 (unknown) • Fields 06–07 (Hour): not 00–23 or not 99 (unknown) • Fields 08–09 (Minutes): not 00–59 or not 99 (unknown)
Data elements	Project Number, Field 01, Fields 02–09
Reference	Group 16 — Special Projects documentation is now accessible through the DAD/NACRS Abstracting Manual application on CIHI's website.

53 Project 440 — Invalid Value for Date and Time of Arterial Puncture (D1620-157)

Rule

When Special Project 440 Endowre (D1620-157)

54 Project 440 — Invalid Value for First Reperfusion Achieved (D1620-158)

Rule

When Project 440 Endovascular Thrombectomy is recorded, it is mandatory to record Field 16 First Reperfusion Achieved.

Reperfusion is achieved when flow is re-established to the ischemic region for the first time. Note that reperfusion may not be sustained by end of the intervention; this element captures whether initial flow occurred.

Criteria	Description
Selection criteria	Project Number 440 is recorded And Field 16 First Reperfusion Achieved is not equal to Y (Yes), N (No) or 9 (Unknown)
Data elements	Project Number, Field 16
Reference	Group 16 — Special Projects documentation is now accessible through the DAD/NACRS Abstracting Manual application on CIHI's website.

55 Project 440 — Invalid Value for Date and Time of First Reperfusion Achieved When First Reperfusion Achieved Is Y (D1620-159)

Rule

When Special Project 440 Endovascular Thrombectomy is recorded and Field 16 First Reperfusion Achieved is recorded with Y (Yes), Fields 17–22 Date and Time of First Reperfusion Achieved must be a valid date and time or 999999 (unknown).

This field captures the time of the first evidence of flow into the affected ischemic territory.

Criteria	Description
Selection criteria	Project Number 440 is recorded And Field 16 First Reperfusion Achieved is Y (Yes) And 1 or more of the following fields are blank or invalid: • Fields 17–18 (Day): not 01–31 or not 99 (unknown) • Fields 19–20 (Hour): not 00–23 or not 99 (unknown) • Fields 21–22 (Minutes): not 00–59 or not 99 (unknown)
Data elements	Project Number, Field 16, Fields 17–22
Reference	Group 16 — Special Projects documentation is now accessible through the DAD/NACRS Abstracting Manual application on CIHI's website.

56 Project 440 — Invalid Value for Final Endovascular Thrombectomy Reperfusion Outcome Achieved (D1620-160)

Rule

When Project 440 Endovascular Thrombectomy is recorded, it is mandatory to record Field 23 Final Endovascular Thrombectomy Reperfusion Outcome Achieved.

This field captures whether final perfusion was achieved and the grade of perfusion at the conclusion of the endovascular intervention, based on evidence during the final angiography imaging run.

Criteria	Description
Selection criteria	Project Number 440 is recorded And Field 23 Final Endovascular Thrombectomy Reperfusion Outcome Achieved is not equal to Y (Yes), N (No) or 9 (Unknown)
Data elements	Project Number, Field 23
Reference	Group 16 — Special Projects documentation is now accessible through the DAD/NACRS Abstracting Manual application on CIHI's website.

57 Number of Previous Term Deliveries and Number of Pre-Term Deliveries Indicate No Previous Delivery and Diagnosis Code or Caesarean Section Status Attribute Indicate Previous Delivery (D1801-120)

Rule

1. When Diagnosis Code O75.701 *Vaginal Delivery Following Caesarean Section*, or O34.201 *Uterine scar due to previous caesarean section*, or O66.401 *Failed trial of labour* is recorded, BOTH data elements Number of Previous Term Deliveries and Number of Previous Pre-Term Deliveries cannot be 00.
2. When Intervention Code 5.MD.60.^^ *Caesarean section delivery* with a Status Attribute **RA** Repeat, Indicated, Planned, **RB** Repeat, Indicated, Emergent or **RC** Repeat, No indicated, Planned is recorded, BOTH data elements Number of Previous Term Deliveries and Number of Previous Pre-Term Deliveries cannot be 00.

Criteria	Description
Selection criteria	Abstracts where Diagnosis Code O75.701, or O34.201 or O66.401 is recorded AND both data elements Number of Previous Term Deliveries and Number of Pre-Term Deliveries is 00. Abstracts where Intervention Code 5.MD.60.^^ with a Status Attribute of RA, RB, or RC is recorded and both data elements Number of Previous Term Deliveries and Number of Pre-Term Deliveries is 00.
Data elements	Number of Previous Term Deliveries, Number of Previous Pre-Term Deliveries, Diagnosis Code, Intervention Code, Status Attribute
Correct case examples	Number of Previous Pre-Term Deliveries is 00 AND Number of Previous Term Deliveries is 00 AND Diagnosis Code is not O75.701 or O34.201 or O66.401 and 5.MD.60.AA *section delivery, lower segment transverse incision, without instrumentation* Status Attribute is PB (Primary, Indicated, Emergent)
Reference	DAD Abstracting Manual: Group 18 — Reproductive Care

58 Missing Value for Number of Previous Term Deliveries and/or Number of Previous Pre-Term Deliveries for Obstetrics Delivered Cases (D1801-135)

Rule

When an obstetrics delivered code is recorded, it is mandatory to report Number of Previous Term Deliveries and Number of Previous Pre-Term Deliveries in some provinces and territories.

Criteria	Description
Selection criteria	Abstracts from NL, NS, NB, PEI (Prince County Hospital and Queen Elizabeth Hospital) ON, MB, SK, AB, BC, NT, YK, and NU where Diagnosis Code from category O10 to O99 with a sixth digit of 1 or 2 is recorded; OR Diagnosis Code Z37.– (outcome of delivery) is assigned Diagnosis Type M; OR Intervention Code 5.MD.50.^^ or 5.MD.60.^^. AND Data element Number of Previous Term Deliveries and/or Number of Previous Pre-Term Deliveries is blank.
Data elements	Number of Previous Term Deliveries, Number of Previous Pre-Term Deliveries, Diagnosis Code, Diagnosis Type, Intervention Code
Correct case examples	Z37.000 (M) Single live birth, pregnancy resulting from both spontaneous ovulation and conception AND Data element: Number of Previous Pre-Term Deliveries is 00 AND Data element: Number of Previous Term Deliveries is 01
Reference	DAD Abstracting Manual: Group 18 — Reproductive Care

59 Scanner Used to Input Product Number and Cement — Product Number but AIM Code Was Not Enabled (D2010-161)

Rule

AIM (Automatic Identification and Mobility) code is a scanner parameter, enabled by scanning a barcode or series of barcodes in the scanner's product reference manual. The presence of the AIM code ensures that CIHI can appropriately extract the product number. There are 2 main barcode formats, HIBC and GS1, to encode the data in the current prosthesis stickers.

Examples

- HIBC format



When the barcode is scanned, the output will look like this:

]C0+H132005764015511/2215162099277E12

The AIM code is composed of the first 3 digits:]C0

- GS1 format



When the barcode is scanned, the output will look like this:

]C10110603295014577

The AIM code is composed of the first 3 digits:]C1

Criteria	Description
Selection criteria	If the first digit of Product Number/Cement — Product Number is “+”, “H” or “h” OR If the first 2 digits of the Product Number/Cement — Product Number are “01” and the total length of Product Number/Cement — Product Number is greater or equal to 16
Data elements	Product Number, Cement — Product Number
Correct case examples	**Applies to data inputted using a barcode scanner only** HIBC format • Scanned output Product Number:]C0+H132005764015511/2215162099277E12 Lot Number:]C0+H132005764015511/2215162099277E12 GS1 format • Scanned output GTIN:]C10110603295014577
References	CJRR barcode bulletin: https://www.cihi.ca/en/scanning-and-entering-product-barcodes-for-the-canadian-joint-replacement-registry or www.cihi.ca/cjrr HIBC specification: https://www.hibcc.org/publication/view/ansi-hibc-2-3-supplier-labeling-standard/ GS1 general specification https://www.gs1.org/sites/default/files/docs/barcodes/GS1_General_Specifications.pdf

Appendix

Appendix A: Post-procedural disorder codes

This list identifes all post-procedural disorder codes. When a code from this list is assigned, it always requires an external cause code. When the applicable external cause is from Y60–Y84, a Diagnosis Cluster must be applied.

E89.0 Postprocedural hypothyroidism

E89.1 Postprocedural hypoinsulinaemia

E89.2 Postprocedural hypoparathyroidism

E89.3 Postprocedural hypopituitarism

E89.4 Postprocedural ovarian failure

E89.5 Postprocedural testicular hypofunction

E89.6 Postprocedural adrenocortical (-medullary) hypofunction

E89.8 Other postprocedural endocrine and metabolic disorders

E89.9 Postprocedural endocrine and metabolic disorder, unspecifed

G97.0 Cerebrospinal fuid leak from spinal puncture

G97.1 Other reactions to spinal and lumbar puncture

G97.2

G97.2

I97.9 Postprocedural disorder of circulatory system, unspecified

J95.00 Haemorrhage from tracheostomy stoma